# Clean catching midstream urine from pigs: Enhancing studies of urinary tract infection

Kristian Stærk<sup>1,2</sup>, Julie May Jensen<sup>1</sup>, Louise Langhorn<sup>3</sup>, Thomas Emil Andersen<sup>1,2</sup>

1. Department of Clinical Microbiology, Odense University hospital. 2. Dept. of Clinical Research, University of Southern Denmark. 3. Biomedical laboratory, University of Southern Denmark


# Background and purpose


Collecting clean urine from pigs requires sedation and catheterization. This is stressful for the animal and prevents daily collection. The purpose of this study was to develop a method for collecting clean-catch urine and to assess the value of such specimens for microbiological culturing in studies of urinary tract infection.

### Methods and results

We used a total of 41 Landrace x Yorkshire pigs. Clean-catch urine (n=43) was collected in the morning hours in sterile cups (from groups of 4-10 pigs). Urine was cultured and specific gravity was assessed. The results was compared to urine collected by catheter (n=60).

Contamination was essentially negligeable. The urine density was significantly higher for specimens collected by clean catch compared to catheter (P<0.0001). Mean collection time was 57 min (SD:31) (range 20-115 min).






#### Figurtekst.


- A. Procedure
- B. Housing (red area is dunging site)
- C. Urine specific gravity (density)
- D. Colony forming units (CFU)


  Culture of midstream urineDyrkning af

  midtstråle-urin på selektiv plade: fra kontrol

(E) og inficeret gris i seriefortynding (F).







# Take-home message

Urine specimens collected by clean catch are preferred over other methods:

- 3R: Refinement (No sedation), reduction (more samples from the same animals)
- Higher quality specimends: No adverse effects of sedatives, no risk of catheter trauma
- Faster and cheaper: No sedation. Urine from 10 pigs can be collected in less the 60 min



1.021

1.014